



Simulation of Earth-Moon-Mars Environments for the Assessment of Organ Doses

Myung-Hee Y. Kim¹, Nathan A. Schwadron², Lawrence W. Townsend³, and Francis A. Cucinotta⁴

¹Division of Space Life Sciences, Universities Space Research Association, Houston, TX 77058, USA

²University of New Hampshire, Durham, NH 03824, USA

³University of Tennessee, Knoxville, TN 37996, USA

⁴NASA Johnson Space Center, Houston, TX 77058, USA



❖ Simulation of Space Radiation Environment

- GCR in interplanetary space^(1,2) represented by GCR deceleration potential, ϕ .
- Trapped protons⁽³⁾ at low Earth orbit (LEO) and the vertical cutoff rigidities for ISS orbit⁽⁴⁾ and geomagnetic transmission factor for GCR at LEO⁽⁵⁾.
- Statistical prediction of solar particle event (SPE) occurrences⁽⁶⁾ and fluences as a function of mission period by a non-homogeneous Poisson process model fitted to a database of proton fluence measurements of SPEs⁽⁷⁾.
- Calculation of Martian atmospheric shielding thickness at elevation for variable solar activities using Mars Global Surveyor data and the distribution of cosine angle.

❖ Assessment of Organ Doses

- Transport properties of the shielding materials and the astronaut's body tissues: NASA BRYNTRN⁽⁸⁾ and HZETRN/QMSFRG code system^(9,10)
- Spherical shield configurations including spacesuit and equipment room of a spacecraft as 0.3 and 5.0 g/cm² aluminum, respectively.
- Body shielding distribution at the sensitive organs of astronaut of computerized anatomical man (CAM) model⁽¹¹⁾
- Acute radiation response and NASA limits analysis with organ dose estimates for mission scenarios^(12,13,14).

ARR Sickness at Severity Level⁽¹²⁾

Severity Level	Upper Gastrointestinal Distress (UG)	Fatigability and Weakness (FW)
1	No effect	No effect
2	Upset stomach, clammy and sweaty, mouth waters	Somewhat tired with mild weakness
3	Nauseated, considerable sweating, swallows frequently to avoid vomiting	Tired with moderate weakness
4	Vomited once or twice, nauseated, and may vomit again	Very tired and weak
5	Vomited several times, including the dry heaves, severe nauseated, and will soon vomit again	Exhausted with almost no strength


Exposure limit by NASA ⁽¹⁴⁾	Organ dose, Gy-Eq
30-d limit at Skin	1.5
30-d limit at Eye	1.0
30-d limit at BFO	0.25
Minimum BFO dose for ARR	0.5

References



- G.D. Badhwar and P.M. O'Neill, Galactic cosmic radiation model and its applications, *Adv. Space Res.*, vol. 17, no. 2, pp. 7-17, 1996.
- P. M. O'Neill, Badhwar-O'Neill galactic cosmic ray model update based on Advanced Composition Explorer (ACE) energy spectra from 1997 to present, *Adv. Space Res.*, vol. 37, no. 9, pp. 1727-1733, 2006.
- J.I. Vette, The NASA/National Space Science Data Center Trapped Radiation Environmental Model Program (1964-1991). NSSDC/WDC-A-R&91-29, 1991.
- D.F. Smart, M.A. Shea, and E.O. Flückiger, Calculated vertical cutoff rigidities for the International Space Station during magnetically quiet times, SH3.6.28, Proceedings of the 26th International Cosmic Ray Conference, August 17-25, 1999, Salt Lake City, Utah, USA.
- J.W. Wilson, J.E. Nealy, G. De Angelis, F.F. Badavi, C.P. Hugger, F.A. Cucinotta, and M.Y. Kim, Dynamic/anisotropic low Earth orbit environmental models, AIAA 2003-6221, Space 2003.
- Kim, M.Y., Hayat, M.J., Feiveson, A.H., Cucinotta, F.A. Prediction of frequency and exposure level of solar particle events. *Health Phys.*, 97(1), 68-81, July 2009.
- Kim, M.Y., Hayat, M.J., Feiveson, A.H., Cucinotta, F.A. Using high-energy proton fluence to improve risk prediction for consequences of solar particle events. *Adv. Space Res.*, 44, 1428-1432, 2009.
- Cucinotta, F.A., Wilson, J.W., Badavi, F. F. Extension to the BRYNTRN code to monoenergetic light ion beams. Washington DC, NASA Report No. TP-3472, 1994.
- J.W. Wilson, F.F. Badave, F.A. Cucinotta, et al. HZETRN: Description of a free-space ion and nucleon transport and shielding computer program. NASA TP-3495, 1995.
- F.A. Cucinotta, J.W. Wilson, P. Saganti, X. Hu, M.Y. Kim, T. Cleghorn, C. Zeitlin, R.K. Tripathi, Isotopic dependence of GCR fluence behind shielding, *Radiat. Meas.* 41 (2006) 1235-1249.
- M.P. Billings, W.R. Yucker, The computerized anatomical man (CAM) model. Washington DC, NASA CR-134043, 1973.
- S. Hu, M.Y. Kim, G.E. McClellan, F.A. Cucinotta, Modeling the acute health effects of astronauts from exposure to large solar particle events, *Health Phys.*, 96(4), 465-476, April 2009.
- F.A. Cucinotta, M.Y. Kim, V. Willingham, K.A. George, Physical and biological organ dosimetry analysis for International Space Station astronauts, *Radiat. Res.*, 170, 127-138, 2008.
- National Research Council/National Academy of Sciences (NRC/NAS), Committee on the Evaluation of Radiation Shielding for Space Exploration. Managing space radiation risk in the new era of space exploration, the National Academies Press; 2008.

Environmental Parameters for Space Radiation Simulation


GCR Environment Represented by Deceleration Potential ϕ



Average GCR Environment for a Given Mission Period



Geomagnetic Transmission Factor (51.6 deg X 450 km)




Environmental parameters		Mission duration				
		6-Month	12-Month	30-Month*		
GCR deceleration parameter (ϕ), MV	Solar minimum	428	428	428		
	Median solar activity	735	743	773		
	Solar maximum	1100	1100	1100		
SPE fluence (Φ_{30}), p/cm ²	Median level	1.39x10 ⁸	9.74x10 ⁸	5.26x10 ⁹		
	August 1972 SPE	8.10x10 ⁹	8.10x10 ⁹	8.10x10 ⁹		
	Upper 95% level	1.08x10 ¹⁰	4.80x10 ¹⁰	1.35x10 ¹¹		
30-Month	Φ_{30} by each segment		Φ_{30} by fraction to 30-month mission			
Mars	Transits	Mars stay	Total	Transits	Mars stay	Total
Mission	12-month	18-month	30-month	12-month	18-month*	30-month
Median	2.78x10 ⁸	2.03x10 ⁹	2.34x10 ⁹	2.10x10 ⁹	3.16x10 ⁹	5.26x10 ⁹
Upper 95%	2.16x10 ¹⁰	7.50x10 ¹⁰	9.66x10 ¹⁰	5.40x10 ¹⁰	8.10x10 ¹⁰	1.35x10 ¹¹

*Organ dose estimates using SPE fluences proportional to the simulated fluence of 30-month mission.

*Organ dose estimates using SPE fluences proportional to the simulated fluence of 30-month mission.

Simulated Distribution of SPE Fluence at 30 MeV




Vertical Distribution of Mars Atmospheric Properties*

Altitude, km	T, °C	p, kPa	ρ , g/cm ³	Atmospheric shielding thickness, g/cm ²	
				Low density model	High density model
8.0	-41.2	0.34	7.64x10 ⁻⁶	0.14	0.19
6.0	-37.0	0.41	8.98x10 ⁻⁶	3.16	4.34
4.0	-35.0	0.49	1.07x10 ⁻⁵	6.73	9.25
2.0	-33.0	0.58	1.27x10 ⁻⁵	10.97	15.08
0.0	-31.0	0.7	1.50x10 ⁻⁵	16.00	22.00
-2.0	-29.0	0.84	1.78x10 ⁻⁵	19.04	26.17
-4.0	-27.0	1.00	2.12x10 ⁻⁵	22.64	31.13
-6.0	-25.0	1.2	2.52x10 ⁻⁵	26.92	37.01
-8.0	-23.0	1.44	2.99x10 ⁻⁵	32.00	44.00


*Temperature and pressure data based on Mars Global Surveyor

Exposure Limit and Acute Radiation Risk Analysis for Mission Scenarios



Annual Effective Dose Comparison





6-month LEO Mission at 51.6°x400 km




6-month Lunar Mission



30-month Lunar Mission (6-month transit – 18-month Mars stay – 6-month transit)



Mars Orbiter Laser Altimeter (MOLA) Topography - Mars Global Surveyor -



Effective Dose of Males with 20 g/cm² Al Shielding on Martian Topography

